

QUESTION

ANSWER: **C**

When a patient is prescribed a medication, the nurse should first determine if the patient is taking any other medications. This is because many medications can interact with each other, leading to either increased or decreased effectiveness, or even toxicity. In this case, the patient is taking a diuretic, which can interact with the prescribed medication.

The nurse should also assess the patient's vital signs, including blood pressure, heart rate, and respiratory rate. This is because the medication can affect these vital signs, and the nurse needs to be able to identify any changes. In this case, the patient's blood pressure is low, which is a common side effect of the medication.

Finally, the nurse should assess the patient's level of consciousness. This is because the medication can cause dizziness or lightheadedness, which can lead to falls or other injuries. In this case, the patient is alert and oriented, which is a good sign.

ANSWER

The correct answer is C. The nurse should first determine if the patient is taking any other medications. This is because many medications can interact with each other, leading to either increased or decreased effectiveness, or even toxicity. In this case, the patient is taking a diuretic, which can interact with the prescribed medication. The nurse should also assess the patient's vital signs, including blood pressure, heart rate, and respiratory rate. This is because the medication can affect these vital signs, and the nurse needs to be able to identify any changes. In this case, the patient's blood pressure is low, which is a common side effect of the medication. Finally, the nurse should assess the patient's level of consciousness. This is because the medication can cause dizziness or lightheadedness, which can lead to falls or other injuries. In this case, the patient is alert and oriented, which is a good sign.

QUESTION

ANSWER: **C**

QUESTION

ANSWER: **C**

When a patient is prescribed a medication, the nurse should first determine if the patient is taking any other medications. This is because many medications can interact with each other, leading to either increased or decreased effectiveness, or even toxicity. In this case, the patient is taking a diuretic, which can interact with the prescribed medication.

The nurse should also assess the patient's vital signs, including blood pressure, heart rate, and respiratory rate. This is because the medication can affect these vital signs, and the nurse needs to be able to identify any changes. In this case, the patient's blood pressure is low, which is a common side effect of the medication.

Finally, the nurse should assess the patient's level of consciousness. This is because the medication can cause dizziness or lightheadedness, which can lead to falls or other injuries. In this case, the patient is alert and oriented, which is a good sign.

The correct answer is C. The nurse should first determine if the patient is taking any other medications. This is because many medications can interact with each other, leading to either increased or decreased effectiveness, or even toxicity. In this case, the patient is taking a diuretic, which can interact with the prescribed medication. The nurse should also assess the patient's vital signs, including blood pressure, heart rate, and respiratory rate. This is because the medication can affect these vital signs, and the nurse needs to be able to identify any changes. In this case, the patient's blood pressure is low, which is a common side effect of the medication. Finally, the nurse should assess the patient's level of consciousness. This is because the medication can cause dizziness or lightheadedness, which can lead to falls or other injuries. In this case, the patient is alert and oriented, which is a good sign.

ANSWER

The correct answer is C. The nurse should first determine if the patient is taking any other medications. This is because many medications can interact with each other, leading to either increased or decreased effectiveness, or even toxicity. In this case, the patient is taking a diuretic, which can interact with the prescribed medication. The nurse should also assess the patient's vital signs, including blood pressure, heart rate, and respiratory rate. This is because the medication can affect these vital signs, and the nurse needs to be able to identify any changes. In this case, the patient's blood pressure is low, which is a common side effect of the medication. Finally, the nurse should assess the patient's level of consciousness. This is because the medication can cause dizziness or lightheadedness, which can lead to falls or other injuries. In this case, the patient is alert and oriented, which is a good sign.

QUESTION

ANSWER: **C**

When a patient is prescribed a medication, the nurse should first determine if the patient is taking any other medications. This is because many medications can interact with each other, leading to either increased or decreased effectiveness, or even toxicity. In this case, the patient is taking a diuretic, which can interact with the prescribed medication.

The nurse should also assess the patient's vital signs, including blood pressure, heart rate, and respiratory rate. This is because the medication can affect these vital signs, and the nurse needs to be able to identify any changes. In this case, the patient's blood pressure is low, which is a common side effect of the medication. Finally, the nurse should assess the patient's level of consciousness. This is because the medication can cause dizziness or lightheadedness, which can lead to falls or other injuries. In this case, the patient is alert and oriented, which is a good sign.